

# A proof of the polycirculant conjecture

Eric Mwambene,  
 Department of Pure and Applied Mathematics,  
 University of the Western Cape,  
 Bellville, 7535, South Africa  
 emwambene@uwc.ac.za

## Abstract

This paper presents a solution of the polycirculant conjecture which states that every vertex-transitive graph  $G$  has an automorphism that permutes the vertices in cycles of the same length. This is done by identifying vertex-transitive graphs as coset graphs. For a coset graph  $H$ , an equivalence relation  $\sim$  is defined on the vertices of cosets with classes as double cosets of the stabiliser and any other proper subgroup  $A'$  of a transitive group  $A$  of  $G$ . Induced left translations of elements of the subgroup  $A'$  are semi-regular since they preserve these double cosets and acts regularly on each of them. The coset graph is equivalent to  $G$  by a theorem of Sabidussi.

*MSC(2000): 05C25, 20B25*

## 1 Introduction

This paper sets out to prove the polycirculant conjecture. The polycirculant conjecture states that every vertex-transitive finite graph is a polycirculant.

A graph  $G = (V, E)$  is a set  $V$  together with an irreflexive and symmetric binary relation  $E$  defined on  $V$ . The two *arcs*  $(x, y), (y, x) \in E$  are identified

into an edge  $[x, y]$ . All graphs discussed in this paper are loopless, connected and finite.

An automorphism of a graph is a permutation of its vertices which preserves edges. The set of all permutations of a graph  $G$  is denoted by  $\text{Aut } G$  and constitutes a group under composition. A graph  $G$  is vertex-transitive if given any two vertices  $x, y \in V(G)$  there is an automorphism which maps  $x$  to  $y$ . Let  $r \geq 1$  and  $s \geq 2$  be integers. An automorphism  $\alpha$  of a graph is said to be  $(r, s)$ -semiregular if it has  $r$  orbits of length  $s$  and *semiregular* if it is  $(r, s)$ -semiregular for some  $r$  and  $s$ . A graph  $G$  of order  $n$  is called a *circulant* if it has a  $(1, n)$ -semiregular automorphism.

It is easy to see that a graph is a circulant if and only if it is Cayley. A natural relaxation of the concept of a graph being a circulant is the concept of it being polycirculant. In 1981, Dragan Marušić [3] asked if every vertex-transitive graph is a polycirculant. The question was also independently asked by D. Jordan [1].

The main thrust of this note is the following result.

**1 Theorem** *Every vertex-transitive graph is a polycirculant.*

To prove the theorem, we first represent vertex-transitive graphs as coset graphs. It is a classical Sabidussi Representation Theorem[5] that vertex-transitive graphs are coset graphs.

For convenience and to fix notation, we will first present the Sabidussi Representation Theorem and then we give the proof of our main result.

## 2 Main Results

### 2.1 Sabidussi's Representation Theorem

In this section we give a presentation of Sabidussi Representation Theorem that is amenable to proving the polycirculant theorem.

Let  $G$  be a vertex-transitive graph and  $A$  a subgroup of  $\text{Aut } G$  which acts

transitively on  $V(G)$ .

Fix  $u \in V(G)$  (base point) and consider the stabilizer of  $u$  in  $A$ :

$$A_u := \{\alpha \in A : \alpha(u) = u\}.$$

Let  $A/A_u$  be the left cosets of  $A_u$  in  $A$  and  $B := \{\alpha \in A : [u, \alpha(u)] \in E(G)\}$ .

Note that  $B$  is a Cayley set in  $A$ , i.e, the identity is not in  $B$  and if  $b \in B$  then  $b^{-1} \in B$ .

We define a graph  $H$  as follows:

$$V(H) := A/A_u,$$

$$[\alpha A_u, \beta A_u] \in E(H) \iff \text{there exists } v \in \beta A_u \text{ such that } v \in \alpha A_u \sigma, \sigma \in B.$$

**2 Lemma** *With the notation as above,  $H$  is isomorphic to  $G$ .*

The constructed graph  $H$  is called a *coset graph*. In view of Lemma 2 we have that

**3 Theorem** (Sabidussi [5]) *Every vertex-transitive graph is a coset graph.*

## 2.2 Proof of the theorem

Here the notation is as in Section 2.1. Our proof of the theorem utilizes the relationship between Cayley graphs and vertex-transitive graphs. The most used characterization of Cayley graph is that given by Sabidussi. It reads as follows:

**4 Proposition** (Sabidussi [5]) *A graph  $G$  is Cayley if and only if  $\text{Aut } G$  contains a subgroup  $A$  which acts regularly on  $V(G)$ .*

However, for our purpose we present yet another characterization which identifies and uses the fact that Cayley graphs should also be considered as coset graphs.

**5 Proposition** *A graph  $G$  is Cayley if and only if there is a transitive subgroup  $A$  of  $\text{Aut } G$  such that  $A_u$  (the stabilizer of  $u$  in  $A$ ) is normal in  $A$  for any  $u \in V(G)$ .*

**Proof**

By Proposition 4, for a given Cayley graph  $G$ ,  $\text{Aut } G$  contains a subgroup  $A$  which acts regularly on  $V(G)$ . Moreover,  $A_u$  contains only the identity element and therefore is trivially normal in  $A$ .

Suppose that  $G$  has it that  $\text{Aut } G$  contains a transitive subgroup  $A$  such that  $A_u$  is normal in  $A$ . Then the quotient group  $A/A_u$  together with the set

$$C := \{\alpha A_u \in A/A_u : \alpha \in A, [u, \alpha u] \in E(G)\}$$

forms a Cayley graph  $\text{Cay}(A/A_u, C)$  isomorphic to  $G$ . ■

To complete the proof of Theorem 1 we need the following lemma.

**6 Lemma** *Let  $G$  be a graph and  $u \in V(G)$ . Let  $A$  be a transitive subgroup of an automorphism group of a graph  $G$  and  $B := \{\alpha \in A : [u, \alpha u] \in E(G)\}$ . If  $A$  contains a proper subgroup  $A'$  generated by  $B' \subset B$ , then it contains a semi-regular element.*

**Proof**

For the vertex-transitive graph  $G$ , let  $H$  be the quotient graph defined as in Section 2.1 by

$$V(H) = A/A_u,$$

$$[\alpha A_u, \alpha' A_u] \in E(H) \iff \text{for some } \tau \in \alpha A_u, \tau' \in \alpha' A_u \text{ s.t. } \tau\beta = \tau', \beta \in B.$$

We define a relation  $\sim$  on  $V(H)$  by

$$\alpha A_u \sim \alpha' A_u \iff \alpha, \alpha' \in A' \tau A_u,$$

i.e.  $\alpha A_u$  and  $\alpha' A_u$  are related if their union is contained in some double coset  $A' \tau A_u$ .

It is clear that  $\sim$  is an equivalence relation.

We have that for any  $\beta \in A'$  the map  $\tilde{\lambda}_\beta : V(H) \rightarrow V(H)$  given by

$$\tilde{\lambda}_\beta(\alpha A_u) = \beta \alpha A_u$$

is an automorphism of  $H$  that preserves equivalence classes of  $\sim$ . Moreover,  $\Lambda_{A'} := \{\tilde{\lambda}_\beta, \beta \in A'\}$  is regular on each double coset. (We can consider the double coset  $A'\tau A_u$  as a set of left cosets in the form

$$A'\tau A_u = \{\beta\alpha A_u : \beta \in A'\}$$

and so for each  $\beta \in A'$  we have

$$\tilde{\lambda}_\beta(A'\tau A_u) = \beta A'\tau A_u = A'\tau A_u$$

for each double coset.) Hence the map  $\tilde{\lambda}_\beta : V(H) \rightarrow V(H)$  is semi-regular. ■

### Proof of Theorem 1

Let  $G$  be a vertex-transitive graph,  $u \in V(G)$  and  $B := \{\alpha \in \text{Aut } G : [u, \alpha u] \in E(G)\}$ .

If there is an element  $\beta \in B$  such that  $\langle \beta \rangle = \text{Aut } G$  then the stabilizer of  $u$  is normal in  $\text{Aut } G$ , hence by Proposition 5,  $G$  is Cayley (Since  $\text{Aut } G$  is Abelian). We therefore have that  $G$  is polycirculant.

Otherwise,  $\langle \beta \rangle, \beta \in B$  is a proper subgroup of  $\text{Aut } G$  and by Lemma 6,  $\text{Aut } G$  contains a semi-regular element. (By considering the equivalence relation  $\sim$  defining double cosets in the form  $\langle \beta \rangle \alpha A_u, \alpha \in \text{Aut } G$  as equivalence classes.) ■

## References

- [1] D. Jordan, Eine symmetrieeigenschaft von graphen, graphentheorie and ihre anwendungen (Stadt Wehlen, 1988), *Dresdner Reihe Forsch.* **9** (1988), 17-20.
- [2] D. Marušič and R. Scapellato, Permutation groups, vertex-transitive graphs and semiregular automorphisms, *European J. Combin.* **19** (1998) 707-712.
- [3] D. Marušič, On vertex symmetric digraphs, *Discrete Mathematics* **36** (1981), 69-81.

- [4] Eric Mwambene, Representing vertex-transitive graphs as groupoid graphs, *Discrete Mathematics*, submitted.
- [5] Gert Sabidussi, Vertex-transitive graphs, *Monatsh. Math* **68** (1964), 426-438.